Trends in Mathematics
Information Center for Mathematical Sciences
Volume 1, Number 1, September 1998, Pages 37–40

ON REGULAR BAER RINGS

JIN YONG KIM AND JAE KEOL PARK

Abstract. We show that a countable π-regular Baer ring is semiprimary. Actually it is shown that a π-regular Baer ring with only countably many idempotents is semiprimary. This rersult allows us to generalize a theorem of Rangaswamy on regular Baer rings. Examples are provided to illustrate and delimit our results.

R will denote an associative ring with unity, $J(R)$ its Jacobson radical and $\mathbf{I}$(R) its set of idempotent elements. We use $|X|$ and $\mathbf{c}$ to denote the cardinality of a set X and the cardinality of the continuum, respectively.

¿From [4], a ring R is called π-regular if for each element $a \in R$ there exist a positive integer n (depending on a) and an element $x \in R$ such that $a^n = a^n x a^n$. A π-regular ring R for which the n in the above can be taken to be 1 is called *regular*. Recall from [5] that a ring R is *Baer* if the right annihilator of every nonempty subset of R is generated as a right ideal by an idempotent. The study of Baer rings has its roots in functional analysis [5]. A ring R is called *right* (resp. *left*) *PP* if the right (resp. left) annihilator of every element of R is generated as a right (resp. left) ideal by an idempotent. Note that every Baer ring is a right and left PP ring.

In 1950 [4], Kaplansky proved that every orthogonally finite (i.e., no infinite set of pairwise orthogonal idempotents) π-regular ring is semilocal. In 1967 [8], Small proved that every orthogonally finite right PP ring is Baer. As a corollary he obtained that every right perfect right PP ring is semiprimary and left PP. In 1974 [7], Rangaswamy established that a regular Baer ring of cardinality less than $\mathbf{c}$ is semisimple Artinian.

The foregoing results motivated us to ask the following natural question: *Is a countable π-regular Baer ring a semiprimary ring?* In this paper we give a positive answer to this question. More generally we will show that if R is a π-regular Baer ring with $|\mathbf{I}(R)| < \mathbf{c}$, then $R = A \oplus B$ where A is a finite direct sum of division rings A_i, where $|A_i| \geq \mathbf{c}$ and B is a semiprimary ring with $|B| < \mathbf{c}$. Rangaswamy's theorem then becomes an immediate corollary of this result.

Recall from [3, p.210] that a ring R is called an *I-ring* if every nonnil right ideal of R contains a nonzero idempotent. It can be checked easily that π-regular rings are I-rings.

Theorem 1. *Let R be an orthogonally finite right PP ring. Then the following conditions are equivalent:*

(i) *R is an I-ring;*

1991 *Mathematics Subject Classification.* 16E50,16D30.

Key words and phrases. von Neumann regular ring, Baer ring, π-regular ring and orthogonally finite ring.

©1998 Information Center for Mathematical Sciences

(ii) *R is a semiprimary ring;*
(iii) *R is a π-regular ring.*

Proof. Since semiprimary rings are π-regular and π-regular rings are I-rings, we only need to prove (i) $\Rightarrow$(ii). Let R be an orthogonally finite I-ring which is right PP. Then by [4, Theorem 2.1] R is semilocal. So it is enough to show that $J(R)$ is nilpotent, where $J(R)$ is the Jacobson radical of R. Write $R = \sum_{i=1}^{n} e_i R$, where $\{e_1, \cdots, e_n\}$ is a complete set of primitive idempotents. Now $J(R) = e_1 J(R) + \cdots + e_n J(R)$. We claim that each $e_i J(R)$ is nilpotent. Suppose that there exists some $e_i J(R)$ which is not nilpotent. Then $(e_i J(R))^2 \neq 0$ for some i, so $e_i x e_i \neq 0$ for some $x \in J(R)$. Consider the map $f_{e_i x} : e_i R \to e_i R$ defined by $f_{e_i x}(e_i r) = e_i x e_i r$. So $\mathrm{Im}(f_{e_i x})$ is projective. Thus $\mathrm{Ker}(f_{e_i x})$=0 since $e_i R$ is indecomposable and $\mathrm{Im}(f_{e_i x}) \neq 0$. Now $e_i x \in J(R)$. Since R is an I-ring, $J(R)$ is nil! and hence there is an integer $m > 1$ such that $(e_i x)^m = 0$ and $(e_i x)^{m-1} \neq 0$. But $(e_i x)^{m-1} \in \mathrm{Ker}\ (f_{e_i x}) = 0$, which is a contradiction. Thus each $e_i J(R)$ is nilpotent. Consequently $J(R)$ is nilpotent.

Immediately we have the following corollary which is due to Small [8].

Corollary 2. *If R is a right perfect right PP ring, then R is a semiprimary ring.*

Example 3. The condition "orthogonally finite" in Theorem 1 is not superfluous. There exists a Baer ring (hence right PP) which is an I-ring, but not π-regular. Now let $R = \{(a_n)_{n=1}^{\infty} \in \prod \mathbb{Q} \mid a_n \in \mathbb{Z} \text{ eventually}\}$, where $\prod \mathbb{Q}$ is the countably infinite direct product of the rationals $\mathbb{Q}$. Then $\prod \mathbb{Q}$ is the maximal ring of quotients $Q(R)$ of R . Since $Q(R)$ is regular self-injective, it is a Baer ring. Also note that the set of all idempotents of $Q(R)$ is that of R. Now for a nonempty subset X of R, it follows that $r_R(X) = r_{Q(R)}(X) \cap R = eQ(R) \cap R$ for some idempotent $e \in R$. Therefore $r_R(X) = eR$, and hence R is a Baer ring. Next, to show that R is an I-ring, let K be a nonzero ideal of R. Then there is a nonzero element, say $x \in K$ with nonzero k-th coordinate, say x_k for some k. Let $y \in Q(R)$ with the k-th coordinate x_k^{-1} and 0 for the other coordinates. Then $y \in R$ and $xy \in K$ is a nonzero idempotent in R. So R is an I-ring. Finally, let $\alpha = (2, 2, \dots) \in R$. If R is π-regular, then there are a positive integer n and an element $\beta \in R$ such that $\alpha^n = \alpha^n \beta \alpha^n$. So there is an integer m such that $2^n = 2^n m 2^n$, which is a contradiction. Thus the ring R cannot be π-regular.

Recall the following result due to Rangaswamy [7,Theorem 1] : Any countable regular Baer ring is semisimple Artinian. In order to give a nontrivial generalization of this result we have the following preparatory lemma.

Lemma 4. *If R is a Baer ring with $|\mathbf{I}(R)| < \mathbf{c}$, then R is orthogonally finite.*

Proof. See [6, Theorem 2].

Theorem 5. *Let R be a π-regular Baer ring with $|\mathbf{I}(R)| < \mathbf{c}$. Then R is semiprimary. In particular, if R is semiprime, then R is semisimple Artinian.*

Proof. This result is a consequence of Lemma 4 and Theorem 1.

Corollary 6. *Let R be a π-regular Baer ring with $|\mathbf{I}(R)| < \mathbf{c}$. Then*
(i) *$R = A \oplus B$ (ring direct sum) and B is countable;*
(ii) *$A = \oplus_{i=1}^{m} A_i$, where each A_i is a division ring with $|A_i| \geq \mathbf{c}$;*

(iii) $B \cong \begin{pmatrix} B_1 & B_{12} & \cdots & B_{1k} \\ 0 & B_2 & \cdots & B_{2k} \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & B_k \end{pmatrix}$, *where each* B_i *is a simple Artinian ring and each* B_{ij} *is a left* B_i*-right* B_j*-bimodule.*

Furthermore, if R *is* π*-regular Baer ring with* $|\mathbf{I}(R)| < \mathbf{c}$ *such that* R *is an algebra over an uncountable field, then* R *is a finite direct sum of division rings.*

Proof. Since R is semiprimary Baer by Theorem 5, it follows that there is a positive integer n by [1,Theorem 4.4] such that

$$R \cong \begin{pmatrix} R_1 & R_{12} & \cdots & R_{1n} \\ 0 & R_2 & \cdots & R_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & R_n \end{pmatrix}$$

with each R_i is simple Artinian and each R_{ij} is a left R_i-right R_j-bimodule. For $1 \leq i \leq n-1$, if R_i is uncountable, then R_i should be a division ring. In this case, since R_{ij} for $i < j \leq n-1$ is a left vector space over a division ring R_i, it follows that $R_{ij} = 0$ whenever R_i is uncountable. Next, suppose that R_n is uncountable. Then it also should be a division ring. In this case, since R_{in} for $1 \leq i \leq n-1$ are right vector spaces over R_n. Thus each $R_{in} = 0$ whenever R_n is a division ring. Thus we can get desired result.

The following result of Rangaswamy [7] becomes an immediate consequence of Theorem 5 or Corollary 6.

Corollary 7. *Let* R *be a regular Baer ring with* $|R| < \mathbf{c}$. *Then* R *is semisimple Artinian.*

The following examples are provided to delimit our results. Recall that a ring R is called *right* (resp. *left*) *weakly regular* (or *fully idempotent*) if $a \in aRaR$ (resp. $a \in RaRa$) for every $a \in R$. Right and left weakly regular rings are called *weakly regular*. Recall [2] that R is a *quasi-Baer* if the right annihilator of every right ideal of R is generated (as a right ideal) by an idempotent.

By Theorem 5, regular Baer rings with only countably many idempotents are semisimple Artinian. So one may raise the following question: *Is a weakly regular Baer ring or a regular quasi-Baer ring with only countably many idempotents a semisimple Artinian ring?* But the following two examples eliminate these possibilities.

Example 8. There exists a weakly regular Baer ring with only countably many idempotents, but it is not semisimple Artinian. In fact let R be the first Weyl algebra over a field of characteristic zero. Then R is a weakly regular Baer ring with only countably many idempotents, but R is not semisimple Artinian.

Example 9. There exists a regular quasi-Baer ring with only countably many idempotents, but it is not semiprimary. Thus Theorem 5 cannot be

LSL.9999 0 9ranscript-

Then R is prime regular, so R is quasi-Baer. Also note that R is countable. However R is not orthogonally finite and hence R cannot be semiprimary. In this case, note that R cannot be a Baer ring because of Theorem 5.

References

1. G. F. Birkenmeier, H. E. Heatherly, J. Y. Kim and J. K. Park, *Semicentral idempotents and triangular representations*, preprint.
2. W. E. Clark, *Twisted matrix units semigroup algebras*, Duke Math.J. **34** (1967), 417–423.
3. N. Jacobson, *Structure of Rings*, Amer. Math. Soc. Colloq. Publ. **37** (1964).
4. I. Kaplansky, *Topological representation of algebras,II*, Trans. Amer. Math. Soc. **68** (1950), 62–75.
5. I. Kaplansky, *Rings of Operators*, Benjamin, New York (1968).
6. J. Y. Kim and J. K. Park, *When is a regular ring semisimple Artinian?*, Math. Japonica **45** (1997), 311–313.
7. K. M. Rangaswamy, *Regular and Baer rings*, Proc. Amer. Math. Soc. **42** (1974), 354–358.
8. L. W. Small, *Semihereditary rings*, Bull. Amer. Math. Soc. **73** (1967), 656–658.

Department of athematics, Kyung Hee University, Suwon 449–701, South Korea
E-mail address: jykim@nms.kunghee.ac.kr

Department of athematics, Busan National University, Busan 609–735, South Korea
E-mail address: jkpark@hyowon.cc.pusan.ac.kr